




Associations between ethnicity and admission to intensive care among women giving birth: a cohort study

J Jardine,^{a,b}  I Gurol-Urganci,^{a,b}  T Harris,^c J Hawdon,^d D Pasupathy,^{e,f} J van der Meulen,^a 
K Walker,^{a,g} the NMPA project team

^a Faculty of Public Health and Policy, London School of Hygiene and Tropical Medicine, London, UK ^b Centre for Quality Improvement and Clinical Audit, Royal College of Obstetricians and Gynaecologists, London, UK ^c Centre for Reproduction Research, Faculty of Health and Life Sciences, De Montfort University, Leicester, UK ^d Royal Free London NHS Foundation Trust, London, UK ^e Department of Women and Children's Health, King's College London, St Thomas's Hospital, London, UK ^f Faculty of Medicine and Health, Westmead Clinical School, University of Sydney, Sydney, NSW, Australia ^g Clinical Effectiveness Unit, Royal College of Surgeons, London, UK

Correspondence: J Jardine, Faculty of Public Health and Policy, London School of Hygiene and Tropical Medicine, 15-17 Tavistock Place, London WC1H 9SH, UK. Email: jennifer.jardine@lshtm.ac.uk

Accepted 26 May 2021.

Objective To determine the association between ethnic group and likelihood of admission to intensive care in pregnancy and the postnatal period.

Design Cohort study.

Setting Maternity and intensive care units in England and Wales.

Population or sample A total of 631 851 women who had a record of a registerable birth between 1 April 2015 and 31 March 2016 in a database used for national audit.

Methods Logistic regression analyses of linked maternity and intensive care records, with multiple imputation to account for missing data.

Main outcome measures Admission to intensive care in pregnancy or postnatal period to 6 weeks after birth.

Results In all, 2.24 per 1000 maternities were associated with intensive care admission. Black women were more than twice as likely as women from other ethnic groups to be admitted (odds

ratio [OR] 2.21, 95% CI 1.82–2.68). This association was only partially explained by demographic, lifestyle, pregnancy and birth factors (adjusted OR 1.69, 95% CI 1.37–2.09). A higher proportion of intensive care admissions in Black women were for obstetric haemorrhage than in women from other ethnic groups.

Conclusions Black women have an increased risk of intensive care admission that cannot be explained by demographic, health, lifestyle, pregnancy and birth factors. Clinical and policy intervention should focus on the early identification and management of severe illness, particularly obstetric haemorrhage, in Black women, in order to reduce inequalities in intensive care admission.

Keywords ethnicity, obstetric haemorrhage, severe maternal morbidity.

Tweetable abstract Black women are almost twice as likely as White women to be admitted to intensive care during pregnancy and the postpartum period; this risk remains after accounting for demographic, health, lifestyle, pregnancy and birth factors.

Please cite this paper as: Jardine J, Gurol-Urganci I, Harris T, Hawdon J, Pasupathy D, van der Meulen J, Walker K; the NMPA project team. Associations between ethnicity and admission to intensive care among women giving birth: a cohort study. BJOG 2021; <https://doi.org/10.1111/1471-0528.16891>.

Introduction

Intensive care admission signifies severe illness requiring additional care and monitoring, with a high risk of mortality. In pregnancy and birth, there are additional short-term and long-term consequences: during pregnancy, severe illness is associated with problems with fetal growth and development, and preterm birth; postnatal admissions frequently result in separation of the mother and baby, with

associated impacts on breastfeeding rates and maternal mental health.¹ Admission to intensive care is considered a marker of severe maternal morbidity.^{2,3}

Women from ethnic minority groups suffer poorer outcomes than women from White ethnic groups during pregnancy and birth in the UK.^{4–7} In the triennium 2016–18, Black women were over four times more likely to die in pregnancy and childbirth than White women.⁸ This is similar to the inequalities that exist in other high-income

countries.^{2,3,9–13} It is unclear to what extent this observed association is explained by differences between ethnic groups in demographic, lifestyle, pregnancy and birth factors, including co-morbidities such as gestational diabetes and hypertension, which are more common in women of ethnic minority backgrounds.^{14,15} The extent to which intensive care admissions in pregnancy and birth vary by country of origin has been examined in cohorts from the Netherlands¹⁶ and Canada;¹⁷ in both countries, migrant women were more likely to have admissions to intensive care. Variation by ethnic group has been examined in the USA,¹⁸ where Black women are more likely to be admitted. No study has previously examined ethnic variation in the UK. Investigating variation in intensive care admission may offer useful insights into potential mechanisms for addressing ethnic inequalities in maternal morbidity and mortality.^{2,3}

This study uses linked maternity and intensive care data from England and Wales, collected for the purposes of national audits, to evaluate the relationship between maternal ethnicity and admissions to intensive care.^{19–21} Routinely collected healthcare data sources offer efficient access to large population samples and the opportunity to examine uncommon outcomes such as admission to intensive care and any associations with maternal demographics or characteristics.

The aims of this study were: (1) to quantify the association between ethnicity and severe morbidity requiring admission to intensive care in pregnancy and the 6 weeks following birth; (2) to understand how this association is explained by adjustment for demographic, lifestyle, pregnancy and birth characteristics and (3) to understand the reasons for maternal admission to critical care among different ethnic groups.

Methods

Data sources

We used a national maternity data set that was linked to hospital admission data for the purposes of a national audit.²¹ This included data routinely collected in the course of clinical care, which was extracted from the maternity information systems (MIS) used in National Health Service (NHS) hospitals in England and Wales. In England, MIS data were linked at patient level using the mother's and baby's dates of birth, NHS numbers and postcodes to records from the Hospital Episode Statistics (HES), an administrative database containing records of all admissions to English NHS hospitals. Linkage was performed using a deterministic algorithm by a trusted third party (NHS Digital). In Wales, data from MIS are collated to form the Maternity Indicators data set (known as MIDs). This was linked at patient level using NHS numbers and

dates of birth to the Patient Episodes Database for Wales (PEDW), an administrative data set by the National Welsh Informatics Service. Details of linkage processes are available elsewhere.²¹ The linked data contained information on births between the 1 April 2015 and 31 March 2016 in five of six boards in Wales and 128 of 134 trusts in England with an obstetric unit.²¹

The maternity data set was also linked to the Intensive Care National Audit and Research Centre (ICNARC) Case Mix Programme Data set. ICNARC routinely collects information on all admissions to adult general intensive care units in England, Wales and Northern Ireland, together with some specialist intensive care units. The ICNARC Case Mix Programme data set contains information about the source, type and reason for admission, and observations, diagnoses and procedures that occur within the intensive care unit.^{19,22} Maternal identifiers (NHS number, date of birth and postcode) for women who gave birth in England and Wales were used by ICNARC to supply records matching all or some of these identifiers for women admitted to intensive care in England and Wales up to 31 March 2017. Further details about the linkage process are available.²²

Definition of variables

Ethnicity was primarily derived from the hospital admission record (HES/PEDW) and infilled where not useable (unknown [ethnos codes 9, X, Z] or missing) from the MIS record. Ethnicity was categorised into groups: White, Asian or British Asian, Black or Black British, Mixed, Other and Unknown or missing. Ethnicity is self-reported to midwives at the time of booking pregnancy and is well, and generally consistently, recorded in hospital data in England at the level of these groups; there are inconsistencies between more granular classifications (e.g. Black African, Black Caribbean may be coded interchangeably).²³

A woman was defined as having an intensive care admission if she had one or more recorded admissions to an intensive care unit in the ICNARC data set within the time frame of estimated date of conception to 6 weeks after birth. The plausible date of conception was calculated as the date of birth plus 14 days minus the gestation in days at birth. A woman was recorded as having a level 3 admission if her admitting or discharging level of care was level 3 (i.e. requiring ventilation support, or with multi-organ failure).

Demographic factors included maternal age and socio-economic status. Maternal age was grouped into six categories (16–24, 25–29, 30–34, 35–39, 40–44, 45 or older). Wider age-bands were used for women under 25 and over 44 years because of the small numbers of women admitted to intensive care at these ages. Socio-economic status was identified using the index of multiple deprivation of the

woman's postcode at the time of birth in England and the postcode of her GP surgery in Wales. The index of multiple deprivation is an area-level ranking of relative deprivation that incorporates information about income, education, employment, crime and the living environment for each of the 32 844 lower super output areas in England and 1909 areas in Wales used for population analysis.²⁴ Using these rankings, areas were separated into population quintiles of relative deprivation.²⁵

Obstetric history included parity (with parity of three or more handled as a single category) and previous caesarean section. Lifestyle factors included maternal body mass index (BMI) and smoking status recorded in MIS at the time of booking the pregnancy. BMI was handled using WHO categories.²⁶

Pregnancy and birth factors included: mode of birth (unassisted vaginal, instrumental vaginal or caesarean section); preterm birth (occurring before 37 weeks of gestation), multiple birth (twins or higher-order multiple) and stillbirth.

Maternal health conditions complicating pregnancy were identified using the International Classification of Diseases, tenth revision codes²⁷ recorded in HES/PEDW in the birth episode. These included diabetes (pre-existing and gestational, handled together because of the low frequency of pre-existing diabetes), pre-eclampsia, pre-existing or gestational hypertension, and placental conditions of morbidly adherent placenta or abruption.

Details of all coding frameworks used are available in Table S1.

Analysis

The primary outcome of interest was admission to an intensive care unit during pregnancy, birth and the postnatal period up to 6 weeks after birth.

To estimate crude odds ratios between ethnic group and intensive care admission, univariate logistic regression models were used. To investigate possible explanations for associations, a series of multivariable logistic regression models with robust estimates of standard errors to account for clustering within hospitals were used to estimate adjusted odds ratios. The first model adjusted for demographic factors: maternal age, ethnic and socio-economic group. The second added the woman's obstetric history (parity and whether she had a previous caesarean section) and lifestyle factors that were present at the onset of pregnancy (BMI and smoking status). The third, 'full' model additionally incorporated health conditions (diabetes, pre-eclampsia, hypertension, cardiac conditions and placental conditions) and pregnancy and birth factors (multiplicity, mode of birth, preterm birth and stillbirth).

Thresholds for admission to intensive care are known to vary with the provision of enhanced care for critically

unwell women within maternity services, as some units provide higher-level care within maternity units and only admissions to critical care units are captured in ICNARC.^{20,22,28} However, care requiring ventilation and for multi-organ failure (level 3) is provided only in intensive care units. For this reason, a sensitivity analysis was carried out using level 3 admission as the outcome in the fully adjusted model.

Levels of missing data were low (<4%) for the majority of variables included in the analysis. However, 6% of women's records were missing information about postcode, which was used to identify socio-economic status, 12% were missing information about ethnicity in both data sources, and 23% were missing information about each of smoking status and BMI at the time of booking. In the regression analyses, multiple imputation using chained equations was used to handle missing values, with regression coefficients estimated using ten imputed data sets and pooled using Rubin's rules.²⁹ Variables used in the imputed data sets included all variables in the multivariable regressions, and also the year of birth and the hospital in which the woman gave birth. Multiple imputation requires the assumption that data are missing at random given the variables used in the imputation model, which may not be met, in particular for ethnicity, smoking status or BMI. To test the sensitivity of findings to these assumptions, the fully adjusted analysis was repeated using only those records with complete information; this has been found to be robust to a wider range of missingness assumptions.³⁰

Primary reasons for admission were available from the intensive care record and were grouped into those directly related to pregnancy and birth and those indirectly related to pregnancy and birth, following a system used for classifying maternal death.^{7,8,22} Details of this classification are available in Table S2. The proportions admitted for each group of reasons were presented by ethnic group.

All analyses were performed in STATA version 14.1 (Stata-Corp, College Station, TX, USA).

Results

A total of 631 851 women were included in the linked data set, of whom 1414 were recorded as being admitted to intensive care during pregnancy, birth and the postnatal period up to 6 weeks, a rate of 2.24 per 1000 maternities. These women each had at least one and a maximum of three recorded admissions to intensive care, with a total of 1619 admissions overall; 261 women (18.5%) had their first admission to intensive care before birth. Of the women admitted to intensive care, 22.3% were recorded as being from ethnic minority groups. (Table 1, Figure S1).

Women were more than twice as likely to be admitted to intensive care if their recorded ethnicity was Black (4.7

Table 1. Characteristics of 631 851 women who gave birth in England and Wales in 2015/16, and 1414 of those women with recorded admissions to intensive care in pregnancy or the postpartum period up to 6 weeks

Risk factor	All women		Women admitted to intensive care	Rate (per 1000 maternities) admitted	Risk factor	All women		Women admitted to intensive care	Rate (per 1000 maternities) admitted
	n	%				n	%		
All	631 851		1 414	2.24					
Ethnic origin					Smoking status				
White	434 297	77.7	931	2.14	Non-smoker	417 542	85.6	923	2.21
Asian	63 795	11.4	147	2.30	Smoker	70 078	14.4	182	2.60
Black	26 900	4.8	125	4.65	Missing	144 231	22.8	309	2.14
Mixed	10 078	1.8	19	1.89	Previous caesarean section	87 501	14.3	347	3.97
Other	23 763	4.3	54	2.27	Missing	20 149	3.2	40	1.99
Missing	73 018	11.6	138	1.89	Recorded diagnoses				
Age group (years)					Hypertension	3208	0.5	28	8.73
Under 25	115 669	18.9	270	2.33	Placental factors	5917	0.9	143	24.17
25–29	174 440	28.6	297	1.70	Pre-eclampsia	11 484	1.8	188	16.37
30–34	190 075	31.1	413	2.17	Cardiac conditions	2036	0.3	67	32.91
35–39	105 849	17.3	298	2.82	Diabetes	32 706	5.2	143	4.37
40–44	23 340	3.8	92	3.94	Gestation				
45 or older	1667	0.3	15	9.00	Term	565 436	92.9	865	1.53
Missing	20 811	3.3	29	1.39	Preterm	42 889	7.1	492	11.47
Socio-economic deprivation (quintile)					Missing	23 526	3.7	57	2.42
Least deprived (1)	99 438	16.8	210	2.11	Multiplicity				
2	84 112	14.2	173	2.06	Singleton birth	613 669	97.1	1 317	2.15
3	112 183	18.9	236	2.10	Multiple birth	18 182	2.9	97	5.33
4	134 759	22.8	294	2.18	Fetal outcome				
Most deprived (5)	161 850	27.3	396	2.45	Livebirth	628 818	99.5	1 345	2.14
Missing	39 509	6.3	105	2.66	Stillbirth	3 033	0.5	69	22.75
Body Mass Index (kg/m ²)					Mode of birth				
<18.5	14 347	2.9	32	2.23	Unassisted vaginal	380 772	61.6	328	0.86
18.5–24.9	236 456	48.4	457	1.93	Instrumental	75 280	12.2	115	1.52
25.0–29.9	131 161	26.8	295	2.25	Caesarean section	161 665	26.2	951	5.88
30.0–34.9	67 672	13.8	163	2.41	Missing	14 134	2.2	20	1.42
35.0–39.9	25 832	5.2	81	3.14	Parity				
≥40.0	13 447	2.8	62	4.61	0	264 133	42.7	621	2.35
Missing	142936	22.6	324	2.27	1	214 572	34.7	396	1.85
Parity					2	86 037	13.9	189	2.20
0	264 133	42.7	621	2.35	3 or more	53 208	8.6	175	3.29
1	214 572	34.7	396	1.85	Missing	13 901	2.2	33	2.37
2	86 037	13.9	189	2.20					
3 or more	53 208	8.6	175	3.29					
Missing	13 901	2.2	33	2.37					

per 1000 maternities) than White (2.1 per 1000 maternities; crude odds ratio [OR] for Black women compared with White women, 2.21, 95% CI 1.82–2.68) but no difference was observed if the recorded ethnicity was Asian (2.3 per 1000), Mixed (1.9 per 1000) or Other (2.3 per 1000) (Tables 1 and 2).

We sought to understand the extent to which adjustment for various characteristics and risk factors could explain the higher intensive care admissions for Black women compared with White women. This was explored using three different models: the first of which adjusted for demographic factors, the second additionally for obstetric history

Table 2. Maternal and pregnancy characteristics associated with admission to intensive care during pregnancy and the early postpartum period up to 6 weeks among women who gave birth in England and Wales in 2015/16

Characteristic	Crude OR	Model 1* (Demographic)		Model 2* (Lifestyle, history)		Model 3* (Pregnancy and birth)	
		Adjusted OR (95% CI)	P value**	Adjusted OR (95% CI)	P value**	Adjusted OR (95% CI)	P value**
Ethnic origin							
White	Ref	Ref	<0.001	Ref	<0.001	Ref	<0.001
Asian	1.08 (0.91–1.28)	1.06 (0.89–1.27)		1.12 (0.94–1.34)		0.98 (0.81–1.19)	
Black	2.21 (1.82–2.68)	2.02 (1.65–2.48)		1.94 (1.57–2.41)		1.69 (1.37–2.09)	
Mixed	0.85 (0.54–1.35)	0.83 (0.52–1.32)		0.84 (0.53–1.33)		0.83 (0.52–1.33)	
Other	1.04 (0.79–1.36)	1.00 (0.76–1.32)		1.06 (0.80–1.40)		1.07 (0.79–1.43)	
Age group (years)							
Under 25	1.37 (1.16–1.62)	1.38 (1.17–1.63)	<0.001	1.35 (1.14–1.60)	<0.001	1.52 (1.27–1.82)	<0.001
25–29	Ref	Ref		Ref		Ref	
30–34	1.27 (1.09–1.47)	1.29 (1.11–1.50)		1.28 (1.10–1.49)		1.15 (0.99–1.34)	
35–39	1.64 (1.40–1.93)	1.66 (1.41–1.95)		1.61 (1.36–1.90)		1.28 (1.08–1.51)	
40–44	2.31 (1.82–2.92)	2.26 (1.78–2.86)		2.07 (1.62–2.64)		1.31 (1.01–1.70)	
45 or older	5.35 (3.17–9.04)	4.89 (2.89–8.27)		4.39 (2.59–7.47)		2.10 (1.23–3.58)	
Socio-economic deprivation (quintile)							
Least deprived (1)	Ref	Ref	0.44	Ref	0.93	Ref	0.93
2	0.96 (0.79–1.18)	0.98 (0.80–1.20)		0.95 (0.78–1.17)		0.95 (0.77–1.17)	
3	0.99 (0.82–1.20)	1.01 (0.84–1.22)		0.96 (0.79–1.17)		0.93 (0.77–1.13)	
4	1.03 (0.86–1.23)	1.03 (0.85–1.24)		0.95 (0.79–1.15)		0.92 (0.76–1.12)	
Most deprived (5)	1.15 (0.98–1.36)	1.14 (0.95–1.37)		1.01 (0.83–1.22)		0.93 (0.77–1.12)	
BMI (kg/m ²)							
<18.5	1.21 (0.87–1.69)			1.22 (0.88–1.69)	<0.001	1.22 (0.84–1.65)	0.006
18.5–24.9	Ref			Ref		Ref	
25.0–29.9	1.16 (1.01–1.33)			1.11 (0.96–1.28)		0.99 (0.85–1.15)	
30.0–34.9	1.26 (1.05–1.52)			1.15 (0.96–1.38)		0.96 (0.79–1.16)	
35.0–39.9	1.71 (1.33–2.19)			1.46 (1.17–1.83)		1.17 (0.90–1.52)	
≥40.0	2.50 (1.91–3.29)			2.10 (1.61–2.75)		1.64 (1.23–2.17)	
Parity							
0	Ref			Ref	<0.001	Ref	0.008
1	0.79 (0.70–0.90)			0.58 (0.51–0.67)		0.95 (0.82–1.19)	
2	0.94 (0.80–1.11)			0.62 (0.52–0.73)		1.05 (0.86–1.26)	
3 or more	1.40 (1.18–1.66)			0.81 (0.68–0.98)		1.33 (1.09–1.61)	
Smoker	1.20 (1.02–1.42)			1.33 (1.13–1.58)	0.001	1.15 (0.95–1.39)	0.14
Previous caesarean section	2.23 (1.98–2.51)			2.41 (2.10–2.76)	<0.001	0.99 (0.85–1.16)	0.92
Maternal conditions							
Diabetes	2.07 (1.74–2.46)					1.26 (1.04–1.53)	0.02
Pre-eclampsia/eclampsia	8.40 (7.20–9.81)					3.11 (2.59–3.74)	<0.001
Hypertension	3.98 (2.74–5.80)					1.59 (1.04–2.42)	0.03
Placental conditions	12.17 (10.22–14.50)					3.46 (2.84–4.22)	<0.001
Cardiac conditions	15.88 (12.37–20.37)					11.28 (8.62–14.77)	<0.001
Mode of birth							
Unassisted vaginal	Ref					Ref	<0.001
Instrumental	1.78 (1.43–2.20)					2.06 (1.65–2.59)	
Caesarean section	6.81 (6.00–7.73)					5.04 (4.31–5.90)	
Fetal complications							
Preterm birth	7.57 (6.78–8.46)					3.53 (3.06–4.06)	<0.001
Multiple birth	4.11 (3.29–5.14)					1.11 (0.86–1.43)	0.41
Stillbirth	10.86 (8.50–13.87)					6.50 (4.86–8.68)	<0.001

*All models are adjusted for variables shown as complete.

**P values for categorical variables are derived using the Wald test.

and lifestyle factors, and the third for these together with pregnancy and birth factors. The increased risk of intensive care admission for Black women was partially explained by adjustment for demographic factors: maternal age and socio-economic status (adjusted OR [aOR] 2.02, 95% CI 1.65–2.48). Lifestyle factors and obstetric history present at the start of pregnancy explained very little of the association (aOR 1.94, 95% CI 1.57–2.41). More of the association was explained by pregnancy and birth characteristics, including presence of co-morbidities, mode of birth, pre-term birth and stillbirth (Table 2). Taking all these factors into account, Black women were 1.7 times more likely to be admitted to intensive care than White women (aOR 1.69, 95% CI 1.37–2.09).

Some complications were associated with particularly high rates of intensive care admission. Following adjustment for demographic, lifestyle, pregnancy and birth factors, women who had pre-eclampsia or placental conditions such as abruption or accreta were three times as likely to be admitted to intensive care (for pre-eclampsia: aOR 3.11, 95% CI 2.59–3.74; for placental conditions: aOR 3.46, 95% CI 2.84–4.22). Women with cardiac conditions were 11 times more likely than women without to be admitted to intensive care (aOR 11.28, 95% CI 8.62–14.77). Women who had a caesarean section were five times as likely (aOR 5.04, 95% CI 4.31–5.90) to be admitted. Women who had a preterm birth were more than three times as likely to be admitted (aOR 3.53, 95% CI

3.06–4.06) and women who had a stillbirth were more than six times as likely (aOR 6.50, 95% CI 4.86–8.68).

These results were robust to a sensitivity analysis restricted to level 3 admissions, although a small increase in risk of intensive care admission in women with diabetes was not apparent in the tighter definition of the outcome. Associations with caesarean birth, placental conditions and stillbirth were stronger with level 3 admission (Table S3). In sensitivity analyses restricted to those women with complete data available (Table S4), the associations with ethnicity were attenuated; this was most evident in the fully adjusted model (Wald *P* value for ethnicity overall 0.09). In these complete case analyses there was much greater uncertainty in the estimates because of the smaller sample size; the adjusted odds ratios for Black ethnicity (in the fully adjusted model, aOR 1.43, 95% CI 1.08–1.90) were within the confidence intervals for the results using imputed data (full model aOR 1.69, 95% CI 1.37–2.09).

Two-thirds (67.1%) of admissions were for a reason directly related to pregnancy, such as obstetric haemorrhage, infection, pre-eclampsia and HELLP syndrome (haemolysis, elevated liver enzymes, low platelets) (Figure 1, Table S5). The proportion of admissions that were due to direct, rather than indirect, reasons, and particularly due to obstetric haemorrhage, was higher among women from Black ethnic origin. Forty-two percent of admissions in Black women were for obstetric haemorrhage compared with 34% in White women. Women with no record of




Figure 1. Reasons for admission by ethnic group, for 1340 admissions with complete ethnic group that occurred in pregnancy or the postpartum period for women who gave birth between 1 April 2015 and 31 March 2016.

ethnic origin were more likely to have an admission for an indirect reason.

Discussion

Main findings

Of women who gave birth in England and Wales in 2015/16, 2.24 per 1000 were admitted to intensive care in pregnancy and the 6 weeks after birth. Black women were more than twice as likely as White women to be admitted. This association was only partially explained by adjustment for demographic, lifestyle, pregnancy and birth characteristics. Women with complications, such as placental factors, pre-eclampsia and stillbirth, were much more likely to be admitted to intensive care. These findings were robust to sensitivity analyses using different definitions of the outcome and methods of handling missing data.

Obstetric haemorrhage accounted for a higher proportion of admissions for Black women than for women from other ethnic group.

Strengths and limitations

The main strengths of this study are its size and design. This is a large cohort study using routinely collected data with a high rate of coverage of births in England and Wales (approximately 92%). The use of electronic patient records, collected for payment purposes, reduces the risk of systematic bias: almost all births and intensive care admissions in the UK occur in the NHS. The ICNARC Case Mix Programme data set for evaluating admissions to intensive care is well established and of high quality.¹⁹ Linkage using identifiers such as NHS numbers ensures that matched records are very likely to be true matches, with women identified as having an intensive care admission being highly likely to have been admitted.

Although the linkage method using NHS number, date of birth and postcode is highly specific, the first limitation is in the potential for missed matches.^{19,20} Although completeness of identifiers is high in both data sets^{19,21} there is no reference standard²² data set to enable evaluation of the linkage quality. This has the potential to cause bias if ethnicity is associated with the likelihood of complete identifiers. In this data set, any bias would be to an under-estimation of effect, as women from ethnic minority groups were less likely to have an NHS number present in the MIS.

Further limitations to this study arise from the missing data within the data set, in particular for ethnicity (12% of records). To account for this, in our primary analyses we use multiple imputation, a methodology that, provided the information about ethnicity is missing at random given all of the other variables in the model, will give unbiased estimates. However, it is possible that this assumption is not

met. It is reassuring that our findings are similar in a complete case analysis, where only those records with complete information about all covariates are included, but in this supplementary analysis the association is substantially attenuated; this may be because the sample size is reduced, or because the true association between Black ethnicity and intensive care admission is smaller than in our primary analysis.

The third limitation is the chosen outcome. Admission to intensive care is considered when a woman is too unwell to be cared for in a maternity unit. The capability of maternity units to provide enhanced or high dependency maternity care varies,^{31,32} therefore the threshold to consider admission may vary between units. It is possible that our findings could be due to systematically lower admission thresholds in hospitals with higher proportions of Black women. However, similar associations were found when the analysis was limited to women requiring care for multi-organ failure or ventilation (Table S3), therapies not provided outside intensive care settings.³³

In our analyses we adjust for factors related to the woman's demographics, lifestyle, pregnancy and birth. In women admitted before the day of birth (18.5% of our population) it is possible that the gestation at birth, mode of birth and stillbirth are causally linked to both ethnic group and the antenatal episode of severe illness indicated by intensive care admission. This can introduce a form of bias where the association is inappropriately attenuated.³⁴ This may partially account for the attenuation of the association between Black ethnicity and likelihood of intensive care admission seen between Model 3 and Model 2.

It may also be that women who were admitted to intensive care differed from those who were not admitted but instead unfortunately died, as the result of a lack of care or escalation as is commonly reported in maternal death.^{7,8,35} Data were not available to us for maternal death that occurred outside the hospital admission in which the woman gave birth, limiting the use of death as an alternative outcome in this study. Any change would be small as maternal death is rare, and any bias would be towards an under-estimation of the effect of ethnicity: Black and Asian women are more likely to die during pregnancy and birth in the UK than White women, with the estimated association larger than that seen in our study.^{7,8}

Interpretation (in light of other evidence)

The overall rate of admission to intensive care during pregnancy and the postnatal period was similar to that reported in other international studies (2–4:1000).^{16,36} Studies from the Netherlands,¹⁶ Canada,¹⁷ and the USA,¹⁸ conducted in local populations, similarly show an association between Black ethnicity or African or Caribbean origin and

admissions to intensive care in pregnancy and the postpartum period. In common with other studies examining severe maternal morbidity in the UK we found no association with socio-economic grouping, reflective of the universal healthcare system.³⁷

Studies from the UK Obstetric Surveillance System^{4,5,37} have demonstrated that women from Black African and Caribbean ethnic groups are more likely to experience severe morbidity, with a similar reported magnitude of effect. The UK Obstetric Surveillance System also found that women from some Asian ethnic groups (Pakistani and Bengali) were more likely to experience severe maternal morbidity, which we did not find.⁴ It is possible that this is masked in our data where we have treated ethnicity in larger groupings to deal with potential coding issues.

The reasons for the association between ethnicity and admission to intensive care or other markers of severe maternal morbidity have been widely hypothesised. Postulated reasons for this association include health at the start of pregnancy, reduced socio-economic status, increased propensity to develop pregnancy-related conditions such as eclampsia, differences in health behaviours, and differences in the way women are treated and listened to during maternity care.^{4,5,38–41} In our study, some of the association between ethnicity and intensive care admission was explained by maternal age and co-morbidity, and by pregnancy and birth factors including caesarean birth, preterm birth, placental conditions and stillbirth. However, even following this adjustment, a substantial association remained. We were unable to account for health behaviours, stress, home environment, experiences of maternal care and aspects of structural inequality that may account for the observed associations.^{42–44}

In this cohort, intensive care admissions for Black women were more commonly due to obstetric haemorrhage than those for women from other ethnic groups. There is a possible biological explanation: Black women are more likely to have leiomyomata or fibroids, benign tumours of the uterine myometrium that prevent the uterus from contracting, which are associated with an increased risk of postpartum haemorrhage.^{45,46} For Black women with increased risk of haemorrhage, appropriate recognition and rapid escalation may avoid the need for additional support and intensive care admission.⁴⁷

A secondary finding of our study was that stillbirth is strongly associated with admission to intensive care. This finding has also been demonstrated in a large study of over 6 million births in California,⁴⁸ which found an increased risk of severe maternal morbidity in women with stillbirth (relative risk 4.77, 95% CI 4.53–5.02). There may be common primary causes leading both to stillbirth and maternal admission to intensive care, such as placental abruption. This requires further study, which was not feasible in this

analysis because information on timing of stillbirth and other events within labour was limited.

Conclusion

Women of Black ethnicity are more than twice as likely as women of other ethnic backgrounds to be admitted to intensive care during pregnancy and birth. Even when demographic, lifestyle, pregnancy and birth characteristics are taken into account, these women are still 1.7 times more likely to be admitted to intensive care.

Further investigation is needed to understand the unexplained increase in risk. Clinical and policy action should focus on the prediction, early identification and management of severe illness and obstetric haemorrhage in Black women to reduce these inequalities. Particular action is also needed to improve monitoring of women with complications including stillbirth, cardiac and placental conditions, given the high risk of intensive care admission in these groups, and to prevent and treat maternal conditions such as hypertension, diabetes and pre-eclampsia. Established procedures, such as the use of early warning scores at regular intervals, should be attentively used in Black women.⁴⁹ If targeted, this has the potential to reduce maternal admissions to intensive care significantly, with an associated reduction in clinical costs and trauma to women and their families.¹

Disclosure of interests

All authors have previously or currently received funding from the Healthcare Quality Improvement Partnership (HQIP) who commissioned the linkage between ICNARC and maternity data for the National Maternity and Perinatal Audit. HQIP had no involvement in the design, analysis or writing of this study. DP additionally is a member of the Ockenden review and the New South Wales Maternal and Perinatal Review Committee.

Contribution to authorship

JJ and KW conceived the study. All authors planned the analysis. JJ conducted the analysis and wrote the first draft of the paper. All authors reviewed and redrafted the study. KW supervised the study.

Details of ethics approval

This study used data routinely collected in clinical care to evaluate service provision and performance and therefore individual consent was not sought. Institutional consent to access the data was provided by the NHS Health Research Authority Confidentiality Advisory Group, approval number 16/CAG/0058. This study was approved by the LSHTM Ethics Committee, approval number 14544, on 4 April 2018.

Funding

The National Maternity and Perinatal Audit is commissioned by the Healthcare Quality Improvement Partnership (HQIP; www.hqip.org.uk) as part of the National Clinical Audit and Patient Outcomes Programme and funded by NHS England and the Scottish and Welsh governments. Neither HQIP nor the funders had any involvement in designing the study; collecting, analysing and interpreting the data; writing the report; or the decision to submit the article for publication.

Patient and public involvement

There was no patient and public involvement in this study.

Acknowledgements

The authors thank Dr Francesca Cavallaro for her comments on a draft of this manuscript. The authors are grateful to the ICNARC team for their provision and linkage of intensive care data from the Case Mix Programme. The authors also thank NHS Digital for their provision of HES data, NHS Trusts in England for the provision of MIS data and the National Welsh Informatics Service for their provision of Welsh maternity and hospital data.

The following individuals are past or current members of the NMPA Project Team: Harriet Aughey, Andrea Blotkamp, Fran Carroll, Megan Coe, George Dunn, Alissa Frémeaux, Rebecca Geary, Ipek Gurol-Urganci, Tina Harris, Jane Hawdon, Jennifer Jardine, Hannah Knight, Lindsey Mamza, Natalie Moitt, Patrick Muller, Dharmindra Pasupathy, Sophie Relph, Louise Thomas, Jan van der Meulen, Lara Waite and Kirstin Webster.

Data availability statement

The data that support the findings of this study are available from ICNARC, NHS Digital, the National Welsh Informatics Service and HQIP. Restrictions apply to the availability of these data, which were used under license for this study. Data are available from the authors with the permission of ICNARC, NHS Digital, the National Welsh Informatics Service and HQIP.

Supporting Information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Figure S1. Flow diagram.

Table S1. Data sources for key variables, together with ICD-10 codes used to define co-morbidities.

Table S2. List of indications for admission and classification system.

Table S3. Sensitivity analysis examining admission to level 3 intensive care among 631 851 women who gave birth in England between 1 April 2015 and 31 March 2016.

Table S4. Sensitivity analysis: complete case analysis.

Table S5. Primary reasons for admission by ethnicity in 1619 admissions among 1414 women who gave birth in England and Wales in 2015/16 and were admitted to intensive care in England during pregnancy and the postpartum period up to 6 weeks (data for Fig 1).

Table S6. Summary characteristics of 631 851 women who gave birth in England and Wales in 2015/16, by ethnic group. ■

References

- 1 Furuta M, Sandall J, Bick D. Women's perceptions and experiences of severe maternal morbidity—a synthesis of qualitative studies using a meta-ethnographic approach. *Midwifery* 2013;30:158–69.
- 2 Main EK, Abreo A, McNulty J, Gilbert W, McNally C, Poeltler D, et al. Measuring severe maternal morbidity: validation of potential measures. *Am J Obstet Gynecol* 2016;214:643.e1–643.e10.
- 3 Roberts CL, Cameron CA, Bell JC, Algert CS, Morris JM. Measuring maternal morbidity in routinely collected health data. *Med Care* 2008;46:786–94.
- 4 Nair M, Kurinczuk JJ, Knight M. Ethnic variations in severe maternal morbidity in the UK—a case control study. *PLoS One* 2014;9:e95086.
- 5 Knight M, Kurinczuk JJ, Spark P, Brocklehurst P. Inequalities in maternal health: national cohort study of ethnic variation in severe maternal morbidities. *BMJ* 2009;338:b542.
- 6 Nair M, Kurinczuk JJ, Knight M. Establishing a national maternal morbidity outcome indicator in England: a population-based study using routine hospital data. *PLoS One* 2016;11:e0153370.
- 7 Knight M, Bunch K, Tuffnell D, Shakespeare J, Kotnis R, Kenyon S, et al. (Eds.). *Saving Lives, Improving Mothers' Care - Lessons learned to inform maternity care from the UK and Ireland Confidential Enquiries into Maternal Deaths and Morbidity 2015-17*. Oxford: National Perinatal Epidemiology Unit, University of Oxford; 2019.
- 8 Knight M, Bunch K, Tuffnell D, Shakespeare J, Kotnis R, Kenyon S, et al. (Eds.). *Saving Lives, Improving Mothers' Care - Lessons learned to inform maternity care from the UK and Ireland Confidential Enquiries into Maternal Deaths and Morbidity 2016-18*. Oxford: National Perinatal Epidemiology Unit, University of Oxford; 2020.
- 9 Howell EA, Egorova NN, Janevic T, Balbierz A, Zeitlin J, Hebert PL. Severe maternal morbidity among hispanic women in New York city: investigation of health disparities. *Obstetrics Gynecol* 2017;129:285–94.
- 10 de Jonge A, van der Goes B, Ravelli A, Amelink-Verburg M, Mol B, Nijhuis J, et al. Perinatal mortality and morbidity in a nationwide cohort of 529,688 low-risk planned home and hospital births. *BJOG* 2009;116:1177–84.
- 11 Jayaratnam S, Burton A, Connan KF, Costa C. Maternal 'near miss' at Royal Darwin Hospital: an analysis of severe maternal morbidity at an Australian regional tertiary maternity unit. *Aust NZJ Obstet Gynaecol* 2016;56:381–6.
- 12 Callaghan WM, Grobman WA, Kilpatrick SJ, Main EK, D'Alton M. Facility-based identification of women with severe maternal morbidity: it is time to start. *Obstetrics Gynecol* 2014;123:978–81.
- 13 Geller SE, Koch AR, Garland CE, MacDonald EJ, Storey F, Lawton B. A global view of severe maternal morbidity: moving beyond maternal mortality. *Reprod Health* 2018;15(Suppl 1):98.

- 14 Bryant AS, Seely EW, Cohen A, Lieberman E. Patterns of pregnancy-related hypertension in black and white women. *Hypertens Pregnancy* 2009;24:281–90.
- 15 Murphy HR, Steel SA, Roland JM, Morris D, Ball V, Campbell PJ, et al. Obstetric and perinatal outcomes in pregnancies complicated by Type 1 and Type 2 diabetes: influences of glycaemic control, obesity and social disadvantage. *Diabetic Med* 2011;28:1060–7.
- 16 Zwart JJ, Dupuis JRO, Richters A, Ory F, van Roosmalen J. Obstetric intensive care unit admission: a 2-year nationwide population-based cohort study. *Intens Care Med* 2009;36:256–63.
- 17 Medcalf KE, Park AL, Vermeulen MJ, Ray JG. Maternal origin and risk of neonatal and maternal ICU admission. *Crit Care Med* 2016;44:1314–26.
- 18 Panchal S, Arria AM, Harris AP. Intensive care utilization during hospital admission for delivery: prevalence, risk factors, and outcomes in a statewide population. *Anesthesiology* 2000;92:1537–44.
- 19 ICNARC. *ICNARC Case Mix Programme: Annual Quality Report 2016/17 for adult critical care* [Internet]. [https://onlinereports.icnarc.org/Reports/2017/12/annual-quality-report-201617-for-adult-critical-care]. Accessed 31 January 2021.
- 20 Harrison DA, Brady AR, Rowan K. Case mix, outcome and length of stay for admissions to adult, general critical care units in England, Wales and Northern Ireland: the Intensive Care National Audit & Research Centre Case Mix Programme Database. *Crit Care* 2004;8:R99.
- 21 NMPA Project Team. *National Maternity and Perinatal Audit Clinical report 2017: revised version*. London, UK: Royal College of Obstetricians and Gynaecologists; 2018.
- 22 Jardine J, NMPA Project Team. *Maternity Admissions to Intensive Care in England, Wales and Scotland in 2015/16*. London, UK: RCOG; 2019.
- 23 Mathur R, Bhaskaran K, Chaturvedi N, Leon DA, vanStaa T, Grundy E, et al. Completeness and usability of ethnicity data in UK-based primary care and hospital databases. *J Public Health* 2013;36:684–92.
- 24 Department for Communities and Local Government. *The English Indices of Deprivation 2015 Statistical Release* [Internet]. [https://www.gov.uk/government/statistics/english-indices-of-deprivation-2015]. Accessed 1 October 2020.
- 25 Abel GA, Barclay ME, Payne RA. Adjusted indices of multiple deprivation to enable comparisons within and between constituent countries of the UK including an illustration using mortality rates. *Bmj Open* 2016;6:e012750.
- 26 World Health Organisation. *Nutrition - Body mass index - BMI* [Internet]. [https://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi]. Accessed 12 August 2020.
- 27 World Health Organisation. *International Statistical Classification of Diseases and Related Health Problems 10th Revision* [Internet]. 2016. [https://icd.who.int/browse10/2016/en]. Accessed 1 May 2020.
- 28 Aoyama K, Pinto R, Ray JG, Hill AD, Scales DC, Lapinsky SE, et al. Variability in intensive care unit admission among pregnant and postpartum women in Canada: a nationwide population-based observational study. *Crit Care* 2019;23:381.
- 29 White IR, Royston P, Wood AM. Multiple imputation using chained equations: Issues and guidance for practice. *Stat Med* 2011;30:377–99.
- 30 Bartlett J, Harel O, Carpenter JR. Asymptotically unbiased estimation of exposure odds ratios in complete records logistic regression. *Am J Epidemiol* 2015;182:730–6.
- 31 Royal College of Anaesthetists/Intensive Care Society. *Care of the critically ill woman in childbirth; enhanced maternal care*. [Internet]; 2018. [https://www.rcoa.ac.uk/sites/default/files/documents/2020-06/EMC-Guidelines2018.pdf] Accessed 2 January 2021.
- 32 Blotkamp A, NMPA Project Team. *National Maternity and Perinatal Audit: Organisational Report 2019* [Internet]. London: RCOG; 2019.
- 33 Intensive Care Society. *Levels of Critical Care for Adult Patients*. 2009.
- 34 Cole SR, Platt RW, Schisterman EF, Chu H, Westreich D, Richardson D, et al. Illustrating bias due to conditioning on a collider. *Int J Epidemiol* 2010;39:417–20.
- 35 Knight M, Bunch K, Cairns A, Cantwell R, Cox P, Kenyon S, et al. *Saving Lives, Improving Mothers' Care Rapid Report: Learning from SARS-CoV-2-related and associated maternal deaths in the UK March – May 2020*. Oxford: National Perinatal Epidemiology Unit, University of Oxford; 2020.
- 36 Zeeman GG. Obstetric critical care: a blueprint for improved outcomes. *Crit Care Med* 2006;34(Suppl):S208–14.
- 37 Lindquist A, Knight M, Kurinczuk JJ. Variation in severe maternal morbidity according to socioeconomic position: a UK national case-control study. *BMJ Open* 2013;3:e002742.
- 38 Somer SJH, Sinkey RG, Bryant AS. Epidemiology of racial/ethnic disparities in severe maternal morbidity and mortality. *Semin Perinatol* 2017;41:258–65.
- 39 Zwart JJ, Jonkers MD, Richters A, Öry F, Bloemenkamp KW, Duvekot JJ, et al. Ethnic disparity in severe acute maternal morbidity: a nationwide cohort study in the Netherlands. *Eur J Public Health* 2011;21:229–34.
- 40 Creanga AA, Bateman BT, Kuklina EV, Callaghan WM. Racial and ethnic disparities in severe maternal morbidity: a multistate analysis, 2008–2010. *Am J Obstet Gynecol* 2014;210:435.e1–435.e8.
- 41 Jonkers M, Richters A, Zwart J, Öry F, van Roosmalen J. Severe maternal morbidity among immigrant women in the Netherlands: patients' perspectives. *Reprod Health Matter* 2011;19:144–53.
- 42 Leimert KB, Olson DM. Racial disparities in pregnancy outcomes: genetics, epigenetics, and allostatic load. *Curr Opin Physiology* 2019;13:155–65.
- 43 Giscombé CL, Lobel M. Explaining disproportionately high rates of adverse birth outcomes among african americans: the impact of stress, racism, and related factors in pregnancy. *Psychol Bull* 2005;131:662–83.
- 44 Raleigh VS, Hussey D, Seccombe I, Hallt K. Ethnic and social inequalities in women's experience of maternity care in England: results of a national survey. *J Roy Soc Med* 2010;103:188–98.
- 45 Peddada SD, Laughlin SK, Miner K, Guyon J-P, Haneke K, Vahdat HL, et al. Growth of uterine leiomyomata among premenopausal black and white women. *Proc National Acad Sci USA* 2008;105:19887–92.
- 46 Longo DL, Bulun SE. Uterine fibroids. *New Engl J Med* 2013;369:1344–55.
- 47 Chandraran E, Krishna A. Diagnosis and management of postpartum haemorrhage. *BMJ* 2017;358: j3875.
- 48 Wall-Wieler E, Carmichael SL, Gibbs RS, Lyell DJ, Girsan AI, El-Sayed YY, et al. Severe maternal morbidity among stillbirth and live birth deliveries in California. *Obstetrics Gynecol* 2019;134:310–7.
- 49 Umar A, Ameh CA, Muriithi F, Mathai M. Early warning systems in obstetrics: a systematic literature review. *PLoS One* 2019;14:e0217864.